

F: Points in Figures -- Rectangles

Given a list of rectangles and a list of points in the x-y plane, determine for each point which figures (if any) contain the point.

Input

There will be N rectangle descriptions, one per line, where $N \leq 10$. The first character will designate the type of figure ("r" for rectangle). This character will be followed by four real values designating the x, y coordinates of the upper left and lower right corners. The end of the list will be signalled by a line containing an asterisk in column 1. The remaining lines will contain the x, y coordinates, one per line, of the points to be tested. The end of this list will be indicated by a point with coordinates 9999.9 9999.9; these values should not be included in the output. Points coinciding with a figure border are not considered inside.

Output

For each point to be tested, write a message of the form:

Point i is contained in figure j

for each figure that contains that point. If the point is not contained in any figure, write a message of the form:

Point i is not contained in any figure

Points and figures should be numbered in the order in which they appear in the input.

Sample Input

```
r 8.5 17.0 25.5 -8.5
r 0.0 10.3 5.5 0.0
r 2.5 12.5 12.5 2.5
*
2.0 2.0
4.7 5.3
6.9 11.2
20.0 20.0
17.6 3.2
5.2 7.8
9999.9 9999.9
```

Sample Output

```
Point 1 is contained in figure 2
Point 2 is contained in figure 2
Point 2 is contained in figure 3
Point 3 is contained in figure 3
Point 4 is not contained in any figure
Point 5 is contained in figure 1
Point 6 is not contained in any figure
```




Diagram of sample input figures and data points